

Spanning trees with small degrees and few leaves<sup>☆</sup>

Eduardo Rivera-Campo

*Departamento de Matemáticas  
 Universidad Autónoma Metropolitana - Iztapalapa  
 Av. San Rafael Atlixco 186, México D.F. 09340, México*

---

**Abstract**

We give an Ore-type condition sufficient for a graph  $G$  to have a spanning tree with small degrees and with few leaves.

*Keywords:*

Spanning tree. Bounded degree. Few leaves.

---

**1. Introduction**

From a classical result by Ore [3] it is well-known that if a simple graph  $G$  with  $n \geq 2$  vertices is such that  $d(u) + d(v) \geq n - 1$  for each pair  $u, v$  of non-adjacent vertices of  $G$ , then  $G$  contains a hamiltonian path.

A *leaf* of a tree  $T$  is a vertex of  $T$  with degree one. A natural generalisation of hamiltonian paths are spanning trees with a small number of leaves. In this direction, Ore's result was generalised by Broersma and Tuinstra [1] to the following theorem.

**Theorem 1.1.** [1] *Let  $s \geq 2$  and  $n \geq 2$  be integers. If  $G$  is a connected simple graph with  $n$  vertices such that  $d(u) + d(v) \geq n - s + 1$ , for each pair  $u, v$  of non-adjacent vertices, then  $G$  contains a spanning tree with at most  $s$  leaves.*

Further related results have been obtained by Egawa *et al* [2] and by Tsugaki and Yamashita [5]. See also [4] for a survey on spanning trees with specific properties.

In this note we consider spanning trees with small degrees as well as with a small number of leaves. Our result is the following.

**Theorem 1.2.** *Let  $n, k$  and  $d_1, d_2, \dots, d_n$  be integers with  $1 \leq k \leq n - 1$  and  $2 \leq d_1 \leq d_2 \leq \dots \leq d_n \leq n - 1$ . If  $G$  is a  $k$ -connected simple graph with vertex set  $V(G) = \{w_1, w_2, \dots, w_n\}$  such that  $d(u) + d(v) \geq n - 1 - \sum_{j=1}^k (d_i - 2)$  for*

---

<sup>☆</sup>Research supported by Conacyt, México

*Email address:* `erc@xanum.uam.mx` (Eduardo Rivera-Campo)

any non-adjacent vertices  $u$  and  $v$  of  $G$ , then  $G$  has a spanning tree  $T$  with at most  $2 + \sum_{j=1}^k (d_j - 2)$  leaves and such that  $d_T(w_j) \leq d_j$  for  $j = 1, 2, \dots, n$ .

## 2. Proof of Theorem 2

Let  $T$  be a largest subtree of  $G$  with at most  $2 + \sum_{j=1}^k (d_j - 2)$  leaves and such that if  $w_j \in V(T)$ , then  $d_T(w_j) \leq d_j$ . Since  $G$  is  $k$ -connected and  $n \geq 2$ , it contains a path with at least  $k+1$  vertices. Therefore, we may assume that tree  $T$  has at least  $k+1$  vertices.

If  $T$  is not a spanning tree, there is a vertex  $w$  of  $G$  not in  $T$ . By Menger's theorem, there are  $k$  internally disjoint paths  $\pi_1, \pi_2, \dots, \pi_k$  in  $G$  joining  $w$  to  $k$  different vertices  $r_1, r_2, \dots, r_k$  of  $T$ .

Let  $n_1$  denote the number of leaves of  $T$ . We claim  $n_1 = 2 + \sum_{j=1}^k (d_j - 2)$ , otherwise there is a vertex  $r_i$  such that  $d_T(r_i) < d_{j_i}$  where  $w_{j_i} = r_i$ . Then  $T' = T \cup \pi_i$  is a subtree of  $G$  with more vertices than  $T$  such that  $d_{T'}(w_j) \leq d_j$  for each  $w_j \in V(T')$  and with at most  $n_1 + 1 \leq 2 + \sum_{j=1}^k (d_j - 2)$  leaves, which contradicts our assumption on the maximality of  $T$ .

Because of Ore's theorem, we can assume  $d_i \geq 3$  for some  $i = 1, 2, \dots, k$ . Since  $T$  has  $n_1 = 2 + \sum_{j=1}^k (d_j - 2) \geq 3$  leaves, as shown above, there is a vertex  $w_j$  of  $T$  such that  $d_T(w_j) \geq 3$ . Suppose there are vertices  $x$  and  $y$  of degree one in  $T$  such that  $xy \in E(G)$ . Since  $T$  is not a path, there is an edge  $zz'$  in the unique  $xy$  path contained in  $T$  with  $d_T(z) \geq 3$ . Let  $T' = (T - zz') + xy$  and notice that  $T'$  is a subtree of  $G$  with  $V(T') = V(T)$ , with less than  $2 + \sum_{j=1}^k (d_j - 2)$  leaves and such that  $d_{T'}(w_j) \leq d_j$  for each  $w_j \in V(T')$ . As above, this is a contradiction and therefore no leaves of  $T$  are adjacent in  $G$ .

Notice that  $d_T(r_1) \geq 2$ , otherwise  $T' = T \cup \pi_1$  would be a tree larger than  $T$ , with the same number of leaves and with  $d_{T'}(w_j) \leq d_j$  for each vertex  $w_j$  of  $T'$ . Let  $u$  and  $v$  be any two leaves of  $T$  with the property that the vertex  $r_1$  lies in the unique  $uv$  path  $T_{uv}$ , contained in  $T$ . Orient the edges of  $T$  in such a way that the corresponding directed tree  $\vec{T}$  is outdirected with root  $u$  (see Fig. 1.)

For each vertex  $z \neq u$  in  $T$  let  $z^-$  be the unique vertex of  $T$  such that  $z^-z$  is an arc of  $\vec{T}$ . Let

$$A = \{y \in V(T) : yv \in E(G)\} \text{ and } B = \{x^- \in V(T) : ux \in E(G)\}.$$

Because of the way the tree  $T$  was chosen, all vertices of  $G$  adjacent to  $u$  or to  $v$  lie in  $T$  and therefore  $|A| = d(v)$ . Let  $x_1$  and  $x_2$  be vertices of  $T$  adjacent to  $u$  in  $G$ , if  $x_1^- = x_2^- = z$  for some vertex  $z$  of  $T$ , let  $T' = (T + ux_1) - zx_1$ . Since  $zx_1$  and  $zx_2$  are edges of  $T$ ,  $d_{T'}(z) \geq 2$  and  $T'$  is a subtree of  $G$  with  $V(T') = V(T)$ ,




Figure 1:  $n = 15$ ,  $k = 4$ ,  $d_1, d_2, \dots, d_{15} = 3$ .

with less than  $2 + \sum_{j=1}^k (d_{i_j} - 2)$  leaves and such that  $d_{T'}(w_j) \leq d_j$  for each  $w_j \in V(T')$ . Again, this is a contradiction, therefore  $|B| = d(u)$ .

Since no vertex in  $A \cup (B \setminus \{u\})$  is a leaf of  $T$ ,

$$|A \cup B| \leq |V(T)| - n_1 + 1 \leq (n - 1) - n_1 + 1 = n - 2 - \sum_{j=1}^k (d_j - 2).$$

Also

$$\begin{aligned} |A \cup B| &= |A| + |B| - |A \cap B| = d(u) + d(v) - |A \cap B| \geq \\ &= n - 1 - \sum_{j=1}^k (d_j - 2) - |A \cap B|. \end{aligned}$$

Therefore  $|A \cap B| \geq 1$ ; let  $z^- \in A \cap B$ . We consider two cases:

*Case 1.* Edge  $z^-z$  lies on the path  $T_{uv}$ .

If  $z = r_1$  (see Fig. 2), let

$$T' = ((T + z^-v) - z^-z) \cup \pi_1 \text{ and}$$




Figure 2:  $T' = ((T + z^-v) - z^-z) \cup \pi_1$

and if  $r_1 \neq z$  (see Fig. 3), let

$$T' = (((T + uz) + z^-v) - r_1^-r_1) - z^-z) \cup \pi_1.$$




Figure 3:  $T' = \left( \left( \left( (T + uz) + z^-v \right) - r_1^- r_1 \right) - z^-z \right) \cup \pi_1$

Both situations lead to a contradiction since  $T'$  is a subtree of  $G$  larger than  $T$ , with at most  $2 + \sum_{j=1}^k (d_j - 2)$  leaves and such that  $d_T(w_j) \leq d_j$  for each  $w_j \in V(T')$ .

*Case 2.* Edge  $z^-z$  does not lie on the path  $T_{uv}$ .

If  $z^-$  lies in  $T_{uv}$ , let  $T'' = (T + uz) - z^-z$  (see Fig. 4).




Figure 4:  $T'' = (T + uz) - z^-z$

And if  $z^-$  does not lie in  $T_{uv}$ , let  $x$  be a vertex in  $T_{uv}$  not in  $T_{uz^-}$  such that  $x^-$  is a vertex in  $T_{uz^-}$  (see Fig. 5). Let

$$T'' = \left( \left( \left( (T + uz) + z^-v \right) - x^-x \right) - z^-z \right)$$

In this case  $T''$  is a subtree of  $G$  with  $V(T'') = V(T)$ , with at most  $n_1 - 1 = 1 + \sum_{j=1}^k (d_j - 2)$  leaves and such that  $d_{T''}(w_j) \leq d_j$  for each  $w_j \in V(T'')$ . As seen above, this is not possible.

Cases 1 and 2 cover all possibilities, therefore  $T$  is a spanning tree of  $G$ .

Let  $k \geq 1$  and  $d_1, d_2, \dots, d_n$  be integers with  $3 \leq d_1 \leq d_2 \leq \dots \leq d_n$  and  $X = \{x_1, x_2, \dots, x_k\}$  and  $Y = \{y_1, y_2, \dots, y_{2-k+d_1+\dots+d_k}\}$  be sets of vertices. The complete bipartite graph  $G$  with bipartition  $(X, Y)$  is  $k$ -connected, has  $n = 2 + \sum_{j=1}^k d_j$  vertices and is such that  $d(u) + d(v) \geq 2k = n - 2 - \sum_{j=1}^k (d_j - 2)$  for any vertices  $u$  and  $v$  of  $G$ . Nevertheless, if  $T$  is a spanning tree of  $G$ , then




Figure 5:  $T'' = (((T + uz) + z^-v) - x^-x) - z^-z$

$d_T(x_j) > d_j$  for some  $j = 1, 2, \dots, k$ . This shows that the condition in Theorem 2 is tight.

## References

- [1] H. Broersma, H. Tuinstra, Independence trees and Hamilton cycles, *J. Graph Theory* **29** (1998), 227 - 237.
- [2] Y. Egawa, H. Matsuda, T. Yamashita, K. Yoshimoto, On a spanning tree with specified leaves, *Graphs Combin.* **24(1)** (2008), 13 - 18.
- [3] O. Ore, Note on hamiltonian circuits, *American Math. Monthly* **67** (1960), 55.
- [4] K. Oseki, T. Yamashita, Spanning trees: A survey, *Graphs Combin.* **27(1)** (2011), 1 - 26.
- [5] M. Tsugaki, T. Yamashita, Spanning trees with few leaves, *Graphs Combin.* **23(5)** (2007), 585 - 598.